

Section 4.4 Conditional Probabilities and Independence

Math 141

Main ideas

Two types of probabilities given events A and B and sample space S:

Probability of A:

$$\Pr(A) = \frac{n(A)}{n(S)}$$

Probability of A, given that B is true (or has occurred):

$$\Pr(A|B) = \frac{n(A \cap B)}{n(B)}$$

$$\Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)} \quad \text{thus} \quad \Pr(A \cap B) = \Pr(B) \Pr(A|B)$$

$$\Pr(B|A) = \frac{\Pr(A \cap B)}{\Pr(A)} \quad \text{thus} \quad \Pr(A \cap B) = \Pr(A) \Pr(B|A)$$

If A is independent of B (and similarly if B is independent of A):

$$\Pr(A|B) = \Pr(A)$$

$$\Pr(A \cap B) = \Pr(B) \cdot \Pr(A|B) = \Pr(B) \cdot \Pr(A) = \Pr(A) \cdot \Pr(B)$$

Observation:

Suppose that A is independent of B. Then $\Pr(A \cap B) = \Pr(A) \cdot \Pr(B|A) = \Pr(A) \cdot \Pr(B)$.
Then $\Pr(B) \cdot \Pr(A|B) = \Pr(A \cap B) = \Pr(A) \cdot \Pr(B)$, which means that $\Pr(A|B) = \Pr(A)$, i.e. B is independent of A. In other words, A and B are independent of each other.

Observations

→


→

→

→

Problems

1. Students in our class: from California (or not) and Female (or not):



	C	C'
F		
F'		

$\Pr(C) =$

$\Pr(C') =$

$\Pr(F) =$

$\Pr(F') =$

$\Pr(C|F) =$

$\Pr(C'|F) =$

$\Pr(C|F') =$

$\Pr(C'|F') =$

$\Pr(F|C) =$

$\Pr(F'|C) =$

$\Pr(F|C') =$

$\Pr(F'|C') =$

2. Roll 2 dice. We're interested in the sum.

E is event that sum is odd.

F is event that sum is ≤ 7 .

$$\Pr(E) =$$

$$\Pr(F) =$$

$$\Pr(E \cap F) =$$

$$\Pr(E|F) =$$

$$\Pr(F|E) =$$

Sum of 2 dice

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

3. Suppose there is a bin with 4 green and 6 blue balls. Choose two balls.

Consider the events: G1 = "first ball is green"

B1 = "first ball card is blue"

G2 = "second ball is green"

B2 = "second ball is blue"

First find:

$$\Pr(\text{both balls are green}) =$$

$$\Pr(\text{both balls are blue}) =$$

$$\Pr(\text{one ball is green, one ball is blue}) =$$

Next find:

$$\Pr(G1) =$$

$$\Pr(B1) =$$

$$\Pr(G2|G1) =$$

$$\Pr(B2|G1) =$$

$$\Pr(G2|B1) =$$

$$\Pr(B2|B1) =$$

$$\Pr(G1 \cap G2) =$$

$$\Pr(B1 \cap B2) =$$

$$\Pr(G1 \cap B2) =$$

$$\Pr(B1 \cap G2) =$$

4. Consider the information in the table at right.

		Political party			Total
		D	R	I	
Gender	M	400	700	300	1400
	F	600	300	200	1100
Total		1000	1000	500	2500

$\Pr(R) =$

$\Pr(F) =$

$\Pr(R|F) =$

$\Pr(F|R) =$

$\Pr(R \cap F) =$

Are “being Republican” and “being female” independent?

What exactly does this mean?

5. Suppose that $\Pr(E) = 0.3$, $\Pr(F) = 0.4$, $\Pr(E \cap F) = 0.2$. Are E and F independent?

$\Pr(E|F) =$

$\Pr(F|E) =$

Also note that $\Pr(E \cap F) \neq \Pr(E) \cdot \Pr(F)$.

6. Suppose that $\Pr(E) = 0.5$, $\Pr(F) = 0.4$, $\Pr(E \cap F) = 0.2$. Are E and F are independent?

$\Pr(E|F) =$

$\Pr(F|E) =$

Also note that $\Pr(E \cap F) = \Pr(E) \cdot \Pr(F)$.

7. A hospital uses two tests to classify blood. Every blood sample is subjected to both tests.
- The first test correctly identifies blood types with probability 0.70.
 - The second test correctly identifies blood type with probability 0.80.
 - The probability that at least one of the tests correctly identifies the blood type is 0.90.
- Find the probability that both tests correctly identify the blood type.

Determine the probability that the first test is correct given that the second test is correct.

Determine the probability that the second test is correct given that the first test is correct.

Are the events “Test 1 correctly identifies the blood type” and “Test 2 correctly identifies the blood type” independent?

8. Suppose we know what percentage of spouses within married couples smoke:

$$\Pr(\text{Woman Smokes}) = .20$$

$$\Pr(\text{Man Smokes}) = .30$$

$$\Pr(\text{Both Smokes}) = .10$$

Is one spouse smoking independent of other smoking, or is there dependency?

$$\Pr(\text{WS} | \text{MS}) =$$


Compare this to $\Pr(\text{WS}) = .20$. So the man smoking makes it more likely that the woman also smokes. The same is true in reverse:

$$\Pr(\text{MS} | \text{WS}) = \quad \text{which} > \Pr(\text{MS}) \text{ by the same proportion.}$$


9. Consider an older married couple. Based on their demographics (age, smoke or not, drink or not, etc.), the probability that the husband will be alive ten years from now is 0.90 and the probability that the wife will be alive ten years from now is 0.95. Assuming that the outcomes (of living and dying) for each are independent, what is the probability that at least one of the two will be alive in ten years?

10. Independence vs. mutually exclusivity:


Independent and mutually exclusive.



Not independent and mutually exclusive.



Independent and not mutually exclusive.



Not independent and not mutually exclusive.

