

Name: \_\_\_\_\_

Problem	T/F	1	2 / 3	4	5	Total
Possible	40	26	14	12	8	100
Received						

**DO NOT OPEN YOUR EXAM UNTIL TOLD TO DO SO.**


**You may use a 3 × 5 card of notes. You will not use a calculator.**

**FOR FULL CREDIT, SHOW ALL WORK RELATED TO FINDING EACH SOLUTION.**

**WARPED** By Mike Cavanaugh



**MUTTS** By Patrick McDonnell



40 points Answer the following 20 True/False questions. Each question is worth 2 points. Note: “True” means *always true* or *necessarily true*. “False” means that it may be true sometimes or under some circumstances, but not always or not necessarily. No explanation is necessary whether true or false.

True False The set of vectors of the form  $\left\{ \begin{bmatrix} a + 1 \\ b \\ 2b \end{bmatrix} \right\}$  (for some  $a$  and  $b$ ) is a subspace of  $R^3$ .

True False For a  $2 \times 2$  matrix, if  $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue 4, then  $\begin{bmatrix} -15 \\ -5 \end{bmatrix}$  is also an eigenvector with eigenvalue  $-20$ .

True False Let  $A$  be a  $5 \times 5$  matrix with just two different eigenvalues. It is possible for  $A$  to have a complete set of 5 linearly independent eigenvectors.

True False If a  $8 \times 5$  matrix has rank 2, then  $\text{nullity } A = 3$ .

True False For a  $3 \times 7$  matrix  $A$ , since there are 4 more columns than rows, the largest possible dimension of  $\text{Nul } A$  is 4.

True False If  $\det A \neq 0$ , then  $\det(A^{-1}) = \frac{1}{\det A}$ .

True False For  $3 \times 3$  matrix  $A$ , if  $\det(A - \lambda I) = \lambda^3 - \lambda$ , then  $A$  has no inverse.

True False If  $3 \times 3$  matrix  $A$  has eigenvalues of 4, 5 and 6, then the eigenvectors of  $A$  form a basis for  $R^3$ .

True False For  $v_1, v_2, v_3 \in R^4$ ,  $\text{Span}\{v_1, v_2, v_3\}$  is a subspace of  $R^4$ .

True False It is possible that  $\{v_1, v_3\}$  is a basis for  $\text{Span}\{v_1, v_2, v_3\}$ .

True False If  $A\vec{v}_1 = \lambda_1\vec{v}_1$  and  $A\vec{v}_2 = \lambda_2\vec{v}_2$  where  $\vec{v}_1$  and  $\vec{v}_2$  are linearly independent, then it must be that  $\lambda_1 \neq \lambda_2$ .

True False The sum of two eigenvectors of a matrix is an eigenvector of that same matrix.

True False There is a value of  $k$  for which  $\begin{bmatrix} 1 & 2 \\ 3 & k \end{bmatrix}$  has an eigenvalue of 0.

True False For  $3 \times 3$  matrix  $A$ ,  $\det(-2A^T) = -6 \det A$ .

True False If the determinant  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = 3$ , then  $\begin{vmatrix} a + 2c & b + 2d \\ c & d \end{vmatrix} = 3$ .

True False The column space of  $A$  is the set of all vectors that can be written  $A\vec{x}$  for some  $\vec{x}$ .

True False If  $\{\vec{v}_1, \dots, \vec{v}_7\}$  spans  $R^7$ , then  $\{\vec{v}_1, \dots, \vec{v}_7\}$  is linearly independent.

True False If  $A$  is a  $3 \times 7$  matrix, then  $\dim \text{Row } A < \dim \text{Col } A$ .

True False The set of matrices  $\left\{ \begin{bmatrix} a & a - b \\ b & c \end{bmatrix} \right\}$  (for some  $a, b$  and  $c$ ) is a subspace of  $M_{2 \times 2}$  (the set of all  $2 \times 2$  matrices).

True False If  $3 \times 3$  matrix  $A$  has rank 2, then  $A\vec{x} = \vec{b}$  has an infinite number of solutions for some right hand side  $\vec{b}$ , no solution for some other right hand side  $\vec{b}$ , and a unique solution for yet another right hand side  $\vec{b}$ .

26 points 1. Consider the matrix  $A = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$ .

/10 Find the eigenvalues  $\lambda_1$  and  $\lambda_2$  and corresponding eigenvectors  $\vec{v}_1$  and  $\vec{v}_2$  of  $A$ .

/1 Once you found the eigenvalues  $\lambda_1$  and  $\lambda_2$  (and before you actually found the eigenvectors  $\vec{v}_1$  and  $\vec{v}_2$ ), how did you know that  $\vec{v}_1$  and  $\vec{v}_2$  would be linearly independent?

/2 Give two different diagonalizations  $PDP^{-1} = [\vec{v}_1 \ \vec{v}_2] \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} [\vec{v}_1 \ \vec{v}_2]^{-1}$  of  $A$ .

$$\left[ \begin{array}{c} \phantom{P} \\ \phantom{D} \\ \phantom{P^{-1}} \end{array} \right]^{-1} \text{ and } \left[ \begin{array}{c} \phantom{P} \\ \phantom{D} \\ \phantom{P^{-1}} \end{array} \right]^{-1}$$

/5 Let  $B$  be the basis for  $R^2$  formed from eigenvectors  $\vec{v}_1$  and  $\vec{v}_2$ . Where  $\vec{x} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$ , find  $[\vec{x}]_B$ , the co-ordinates of  $\vec{x}$  with respect to  $B$ . (That is, find  $c_1$  and  $c_2$  so that  $\vec{x} = c_1\vec{v}_1 + c_2\vec{v}_2$ .)

/8 Where  $\vec{x} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$ , find  $A^5\vec{x}$ .

5 points 2. Use Cramer's Rule to find  $x_3$  in the linear system

$$\begin{array}{rcl} 2x_1 & & = 2 \\ 3x_1 + 5x_2 & & = -2 \\ 4x_1 & + x_3 & = 2 \end{array}$$

10 points 3. Consider probability matrix  $P = \begin{bmatrix} .6 & .3 \\ .4 & .7 \end{bmatrix}$  and initial vector  $\vec{x}_0 = \begin{bmatrix} 70 \\ 0 \end{bmatrix}$ . Let  $\vec{x}_k = A^k \vec{x}_0$ .

/1 Find  $\vec{x}_1$ .

/5 Find the eigenvector of  $P$  that corresponds to eigenvalue 1.

/2 Find  $P^\infty$ .

/2 Where  $\vec{x}_0 = \begin{bmatrix} 70 \\ 0 \end{bmatrix}$ , find  $\vec{x}_\infty$ .

12 points 4. Consider matrix  $A$ , which is row equivalent to matrix  $B$ :

$$A = \begin{bmatrix} 1 & 1 & -3 & 7 & 9 & -9 \\ 1 & 2 & -4 & 10 & 13 & -12 \\ 1 & -1 & -1 & 1 & 1 & -3 \\ 1 & -3 & 1 & -5 & -7 & 3 \\ 1 & -2 & 0 & 0 & -5 & -4 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & -2 & 0 & 9 & 2 \\ 0 & 1 & -1 & 0 & 7 & 3 \\ 0 & 0 & 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Find the following:

/2  $\text{rank } A =$

/2  $\dim \text{Nul } A =$

/2 A basis for  $\text{Row } A$  :

/2 A basis for  $\text{Col } A$  :

/4 A basis for  $\text{Nul } A$ :

4 points 5. For  $n \times n$  matrix  $A$  with eigenvalue  $\lambda$ , show that the eigenspace

$$\lambda(A) = \{ \vec{x} : A\vec{x} = \lambda\vec{x} \}$$

is a subspace of  $R^n$ .

4 points 6. If the shape on the left is transformed into the shape on right using the transformation  $T(\vec{x}) = A\vec{x}$ , where

$$A = \begin{bmatrix} 1 & 2 \\ -3 & 1 \end{bmatrix},$$

what is the area of the shape on right?

