



Math 260 Homework 4.2


38. [M] The graph of $f(t)$ is given below. A conjecture is that $f(t) = \sin 3t$.



The graph of $g(t)$ is given below. A conjecture is that $g(t) = \cos 4t$.



The graph of $h(t)$ is given below. A conjecture is that $h(t) = \sin 5t$.



4.2 SOLUTIONS

Notes: This section provides a review of Chapter 1 using the new terminology. Linear transformations are introduced quickly since students are already comfortable with the idea from \mathbb{R}^n . The key exercises are 17–26, which are straightforward but help to solidify the notions of null spaces and column spaces. Exercises 30–36 deal with the kernel and range of a linear transformation and are progressively more advanced theoretically. The idea in Exercises 7–14 is for the student to use Theorems 1, 2, or 3 to determine whether a given set is a subspace.

1. One calculates that

$$Aw = \begin{bmatrix} 3 & -5 & -3 \\ 6 & -2 & 0 \\ -8 & 4 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \\ -4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

so w is in $\text{Nul } A$.

2. One calculates that

$$Aw = \begin{bmatrix} 2 & 6 & 4 \\ -3 & 2 & 5 \\ -5 & -4 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

so w is in $\text{Nul } A$.

3. First find the general solution of $Ax = 0$ in terms of the free variables. Since

$$[A \ 0] = \begin{bmatrix} 1 & 0 & -2 & 4 & 0 \\ 0 & 1 & 3 & -2 & 0 \end{bmatrix},$$

the general solution is $x_1 = 2x_3 - 4x_4$, $x_2 = -3x_3 + 2x_4$, with x_3 and x_4 free. So

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 2 \\ 0 \\ 1 \end{bmatrix},$$

and a spanning set for $\text{Nul } A$ is

$$\left\{ \begin{bmatrix} 2 \\ -3 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

4. First find the general solution of $Ax = 0$ in terms of the free variables. Since

$$[A \ 0] = \begin{bmatrix} 1 & -3 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix},$$

the general solution is $x_1 = 3x_2$, $x_3 = 0$, with x_2 and x_4 free. So

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_2 \begin{bmatrix} 3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix},$$

and a spanning set for $\text{Nul } A$ is

$$\left\{ \begin{bmatrix} 3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

5. First find the general solution of $Ax = 0$ in terms of the free variables. Since

$$[A \ 0] = \begin{bmatrix} 1 & -4 & 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & -5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix},$$

the general solution is $x_1 = 4x_2 - 2x_4$, $x_3 = 5x_4$, $x_5 = 0$, with x_2 and x_4 free. So

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_2 \begin{bmatrix} 4 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ 5 \\ 1 \\ 0 \end{bmatrix},$$

and a spanning set for $\text{Nul } A$ is

$$\left\{ \begin{bmatrix} 4 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 5 \\ 1 \\ 0 \end{bmatrix} \right\}.$$

6. First find the general solution of $A\mathbf{x} = \mathbf{0}$ in terms of the free variables. Since

$$[A \ \mathbf{0}] = \begin{bmatrix} 1 & 0 & 5 & -6 & 1 & 0 \\ 0 & 1 & -3 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

the general solution is $x_1 = -5x_3 + 6x_4 - x_5$, $x_2 = 3x_3 - x_4$, with x_3 , x_4 , and x_5 free. So

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_3 \begin{bmatrix} -5 \\ 3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 6 \\ -1 \\ 0 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix},$$

and a spanning set for $\text{Nul } A$ is

$$\left\{ \begin{bmatrix} -5 \\ 3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 6 \\ -1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

7. The set W is a subset of \mathbb{R}^3 . If W were a vector space (under the standard operations in \mathbb{R}^3), then it would be a subspace of \mathbb{R}^3 . But W is not a subspace of \mathbb{R}^3 since the zero vector is not in W . Thus W is not a vector space.

8. The set W is a subset of \mathbb{R}^3 . If W were a vector space (under the standard operations in \mathbb{R}^3), then it would be a subspace of \mathbb{R}^3 . But W is not a subspace of \mathbb{R}^3 since the zero vector is not in W . Thus W is not a vector space.

9. The set W is the set of all solutions to the homogeneous system of equations $p - 3q - 4s = 0$,

$$2p - s - 5r = 0. \text{ Thus } W = \text{Nul } A, \text{ where } A = \begin{bmatrix} 1 & -3 & -4 & 0 \\ 2 & 0 & -1 & -5 \end{bmatrix}. \text{ Thus } W \text{ is a subspace of } \mathbb{R}^4 \text{ by}$$

Theorem 2, and is a vector space.

10. The set W is the set of all solutions to the homogeneous system of equations $3a + b - c = 0$,

$$a + b + 2c - 2d = 0. \text{ Thus } W = \text{Nul } A, \text{ where } A = \begin{bmatrix} 3 & 1 & -1 & 0 \\ 1 & 1 & 2 & -2 \end{bmatrix}. \text{ Thus } W \text{ is a subspace of } \mathbb{R}^4 \text{ by}$$

Theorem 2, and is a vector space.

11. The set W is a subset of \mathbb{R}^4 . If W were a vector space (under the standard operations in \mathbb{R}^4), then it would be a subspace of \mathbb{R}^4 . But W is not a subspace of \mathbb{R}^4 since the zero vector is not in W . Thus W is not a vector space.

12. The set W is a subset of \mathbb{R}^4 . If W were a vector space (under the standard operations in \mathbb{R}^4), then it would be a subspace of \mathbb{R}^4 . But W is not a subspace of \mathbb{R}^4 since the zero vector is not in W . Thus W is not a vector space.

13. An element \mathbf{w} on W may be written as

$$\mathbf{w} = c \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + d \begin{bmatrix} -6 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 & -6 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix}$$

where c and d are any real numbers. So $W = \text{Col } A$ where $A = \begin{bmatrix} 1 & -6 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$. Thus W is a subspace of \mathbb{R}^3

by Theorem 3, and is a vector space.

14. An element \mathbf{w} on W may be written as

$$\mathbf{w} = s \begin{bmatrix} -1 \\ 1 \\ 5 \end{bmatrix} + t \begin{bmatrix} 3 \\ -2 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 & 3 \\ 1 & -2 \\ 5 & -1 \end{bmatrix} \begin{bmatrix} s \\ t \end{bmatrix}$$

where a and b are any real numbers. So $W = \text{Col } A$ where $A = \begin{bmatrix} -1 & 3 \\ 1 & -2 \\ 5 & -1 \end{bmatrix}$. Thus W is a subspace of

\mathbb{R}^3 by Theorem 3, and is a vector space.

15. An element in this set may be written as

$$r \begin{bmatrix} 0 \\ 1 \\ 3 \\ 2 \end{bmatrix} + s \begin{bmatrix} 2 \\ -1 \\ 1 \\ -1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 1 \\ 1 & -1 & 2 \\ 3 & 1 & 0 \\ 2 & -1 & -1 \end{bmatrix} \begin{bmatrix} r \\ s \\ t \end{bmatrix}$$

where r , s and t are any real numbers. So the set is $\text{Col } A$ where $A = \begin{bmatrix} 0 & 2 & 1 \\ 1 & -1 & 2 \\ 3 & 1 & 0 \\ 2 & -1 & -1 \end{bmatrix}$.

16. An element in this set may be written as

$$b \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix} + c \begin{bmatrix} -1 \\ 0 \\ 3 \\ 1 \end{bmatrix} + d \begin{bmatrix} 0 \\ 3 \\ -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & 3 & -3 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} b \\ c \\ d \end{bmatrix}$$

where b , c and d are any real numbers. So the set is $\text{Col } A$ where $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & 3 & -3 \\ 0 & 1 & 1 \end{bmatrix}$.

17. The matrix A is a 4×2 matrix. Thus

- Nul A is a subspace of \mathbb{R}^2 , and
- Col A is a subspace of \mathbb{R}^4 .

18. The matrix A is a 4×3 matrix. Thus

- Nul A is a subspace of \mathbb{R}^3 , and
- Col A is a subspace of \mathbb{R}^4 .

19. The matrix A is a 2×5 matrix. Thus

- Nul A is a subspace of \mathbb{R}^5 , and
- Col A is a subspace of \mathbb{R}^2 .

20. The matrix A is a 1×5 matrix. Thus

- Nul A is a subspace of \mathbb{R}^5 , and
- Col A is a subspace of $\mathbb{R}^1 = \mathbb{R}$.

21. Either column of A is a nonzero vector in Col A . To find a nonzero vector in Nul A , find the general solution of $Ax = \mathbf{0}$ in terms of the free variables. Since

$$[A \quad \mathbf{0}] = \begin{bmatrix} 1 & -2/3 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{ the general solution is } x_1 = (2/3)x_2, \text{ with } x_2 \text{ free. Letting } x_2 \text{ be a}$$

nonzero value (say $x_2 = 3$) gives the nonzero vector

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

which is in Nul A .

22. Any column of A is a nonzero vector in Col A . To find a nonzero vector in Nul A , find the general solution of $Ax = \mathbf{0}$ in terms of the free variables. Since

$$[A \quad \mathbf{0}] = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

the general solution is $x_1 = -x_3$, $x_2 = -x_3$, with x_3 free. Letting x_3 be a nonzero value (say $x_3 = -1$) gives the nonzero vector

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

which is in Nul A .

23. Consider the system with augmented matrix $[A \quad \mathbf{w}]$. Since

$$[A \quad \mathbf{w}] = \begin{bmatrix} 1 & -2 & -1 \\ 0 & 0 & 0 \end{bmatrix},$$

the system is consistent and \mathbf{w} is in Col A . Also, since

$$A\mathbf{w} = \begin{bmatrix} -2 & 4 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

\mathbf{w} is in Nul A .

24. Consider the system with augmented matrix $[A \quad \mathbf{w}]$. Since

$$[A \quad \mathbf{w}] = \begin{bmatrix} 1 & 0 & 0 & -1 & 1 \\ 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

the system is consistent and \mathbf{w} is in Col A . Also, since

$$A\mathbf{w} = \begin{bmatrix} 10 & -8 & -2 & -2 \\ 0 & 2 & 2 & -2 \\ 1 & -1 & 6 & 0 \\ 1 & 1 & 0 & -2 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

\mathbf{w} is in Nul A .

25. a. True. See the definition before Example 1.

b. False. See Theorem 2.

c. True. See the remark just before Example 4.

d. False. The equation $A\mathbf{x} = \mathbf{b}$ must be consistent for every \mathbf{b} . See #7 in the table on page 204.

e. True. See Figure 2.

f. True. See the remark after Theorem 3.

26. a. True. See Theorem 2.

b. True. See Theorem 3.

c. False. See the box after Theorem 3.

d. True. See the paragraph after the definition of a linear transformation.

e. True. See Figure 2.

f. True. See the paragraph before Example 8.

27. Let A be the coefficient matrix of the given homogeneous system of equations. Since $Ax = \mathbf{0}$ for

$$\mathbf{x} = \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}, \mathbf{x} \text{ is in Nul } A. \text{ Since Nul } A \text{ is a subspace of } \mathbb{R}^3, \text{ it is closed under scalar multiplication. Thus}$$

$10\mathbf{x} = \begin{bmatrix} 30 \\ 20 \\ -10 \end{bmatrix}$ is also in Nul A , and $x_1 = 30, x_2 = 20, x_3 = -10$ is also a solution to the system of equations.

28. Let A be the coefficient matrix of the given systems of equations. Since the first system has a

solution, the constant vector $\mathbf{b} = \begin{bmatrix} 0 \\ 1 \\ 9 \end{bmatrix}$ is in Col A . Since Col A is a subspace of \mathbb{R}^3 , it is closed under

scalar multiplication. Thus $5\mathbf{b} = \begin{bmatrix} 0 \\ 5 \\ 45 \end{bmatrix}$ is also in Col A , and the second system of equations must thus have a solution.

29. a. Since $A\mathbf{0} = \mathbf{0}$, the zero vector is in Col A .

b. Since $Ax + A\mathbf{w} = A(\mathbf{x} + \mathbf{w})$, $Ax + A\mathbf{w}$ is in Col A .

c. Since $c(Ax) = A(c\mathbf{x})$, cAx is in Col A .

30. Since $T(\mathbf{0}_W) = \mathbf{0}_W$, the zero vector $\mathbf{0}_W$ of W is in the range of T . Let $T(\mathbf{x})$ and $T(\mathbf{w})$ be typical elements in the range of T . Then since $T(\mathbf{x}) + T(\mathbf{w}) = T(\mathbf{x} + \mathbf{w})$, $T(\mathbf{x}) + T(\mathbf{w})$ is in the range of T and the range of T is closed under vector addition. Let c be any scalar. Then since $cT(\mathbf{x}) = T(c\mathbf{x})$, $cT(\mathbf{x})$ is in the range of T and the range of T is closed under scalar multiplication. Hence the range of T is a subspace of W .

31. a. Let \mathbf{p} and \mathbf{q} be arbitrary polynomials in \mathcal{P}_2 , and let c be any scalar. Then

$$T(\mathbf{p} + \mathbf{q}) = \begin{bmatrix} (\mathbf{p} + \mathbf{q})(0) \\ (\mathbf{p} + \mathbf{q})(1) \end{bmatrix} = \begin{bmatrix} \mathbf{p}(0) + \mathbf{q}(0) \\ \mathbf{p}(1) + \mathbf{q}(1) \end{bmatrix} = \begin{bmatrix} \mathbf{p}(0) \\ \mathbf{p}(1) \end{bmatrix} + \begin{bmatrix} \mathbf{q}(0) \\ \mathbf{q}(1) \end{bmatrix} = T(\mathbf{p}) + T(\mathbf{q})$$

and

$$T(c\mathbf{p}) = \begin{bmatrix} (c\mathbf{p})(0) \\ (c\mathbf{p})(1) \end{bmatrix} = c \begin{bmatrix} \mathbf{p}(0) \\ \mathbf{p}(1) \end{bmatrix} = cT(\mathbf{p})$$

so T is a linear transformation.

b. Any quadratic polynomial \mathbf{q} for which $\mathbf{q}(0) = 0$ and $\mathbf{q}(1) = 0$ will be in the kernel of T . The

polynomial \mathbf{q} must then be a multiple of $\mathbf{p}(t) = t(t-1)$. Given any vector $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ in \mathbb{R}^2 , the

polynomial $\mathbf{p} = x_1 + (x_2 - x_1)t$ has $\mathbf{p}(0) = x_1$ and $\mathbf{p}(1) = x_2$. Thus the range of T is all of \mathbb{R}^2 .

32. Any quadratic polynomial \mathbf{q} for which $\mathbf{q}(0) = 0$ will be in the kernel of T . The polynomial \mathbf{q} must then be $\mathbf{q} = at + bt^2$. Thus the polynomials $\mathbf{p}_1(t) = t$ and $\mathbf{p}_2(t) = t^2$ span the kernel of T . If a vector is in the range of T , it must be of the form $\begin{bmatrix} a \\ a \end{bmatrix}$. If a vector is of this form, it is the image of the

polynomial $\mathbf{p}(t) = a$ in \mathcal{P}_2 . Thus the range of T is $\left\{ \begin{bmatrix} a \\ a \end{bmatrix} : a \text{ real} \right\}$.

33. a. For any A and B in $M_{2 \times 2}$ and for any scalar c ,

$$T(A+B) = (A+B) + (A+B)^T = A+B + A^T + B^T = (A+A^T) + (B+B^T) = T(A) + T(B)$$

and

$$T(cA) = (cA)^T = c(A^T) = cT(A)$$

so T is a linear transformation.

b. Let B be an element of $M_{2 \times 2}$ with $B^T = B$, and let $A = \frac{1}{2}B$. Then

$$T(A) = A + A^T = \frac{1}{2}B + \left(\frac{1}{2}B\right)^T = \frac{1}{2}B + \frac{1}{2}B^T = \frac{1}{2}B + \frac{1}{2}B = B$$

c. Part b. showed that the range of T contains the set of all B in $M_{2 \times 2}$ with $B^T = B$. It must also be shown that any B in the range of T has this property. Let B be in the range of T . Then $B = T(A)$ for some A in $M_{2 \times 2}$. Then $B = A + A^T$, and

$$B^T = (A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T = B$$

so B has the property that $B^T = B$.

d. Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be in the kernel of T . Then $T(A) = A + A^T = \mathbf{0}$, so

$$A + A^T = \begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} a & c \\ b & d \end{bmatrix} = \begin{bmatrix} 2a & c+b \\ b+c & 2d \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Solving it is found that $a = d = 0$ and $c = -b$. Thus the kernel of T is $\left\{ \begin{bmatrix} 0 & b \\ -b & 0 \end{bmatrix} : b \text{ real} \right\}$.

34. Let \mathbf{f} and \mathbf{g} be any elements in $C[0, 1]$ and let c be any scalar. Then $T(\mathbf{f})$ is the antiderivative \mathbf{F} of \mathbf{f} with $\mathbf{F}(0) = 0$ and $T(\mathbf{g})$ is the antiderivative \mathbf{G} of \mathbf{g} with $\mathbf{G}(0) = 0$. By the rules for antidifferentiation $\mathbf{F} + \mathbf{G}$ will be an antiderivative of $\mathbf{f} + \mathbf{g}$, and $(\mathbf{F} + \mathbf{G})(0) = \mathbf{F}(0) + \mathbf{G}(0) = 0 + 0 = 0$. Thus $T(\mathbf{f} + \mathbf{g}) = T(\mathbf{f}) + T(\mathbf{g})$. Likewise $c\mathbf{f}$ will be an antiderivative of $c\mathbf{f}$, and $(c\mathbf{F})(0) = c\mathbf{F}(0) = c \cdot 0 = 0$. Thus $T(c\mathbf{f}) = cT(\mathbf{f})$, and T is a linear transformation. To find the kernel of T , we must find all functions f in $C[0, 1]$ with antiderivative equal to the zero function. The only function with this property is the zero function $\mathbf{0}$, so the kernel of T is $\{\mathbf{0}\}$.

35. Since U is a subspace of V , $\mathbf{0}_V$ is in U . Since T is linear, $T(\mathbf{0}_V) = \mathbf{0}_W$. So $\mathbf{0}_W$ is in $T(U)$. Let $T(\mathbf{x})$ and $T(\mathbf{y})$ be typical elements in $T(U)$. Then \mathbf{x} and \mathbf{y} are in U , and since U is a subspace of V , $\mathbf{x} + \mathbf{y}$ is also in U . Since T is linear, $T(\mathbf{x}) + T(\mathbf{y}) = T(\mathbf{x} + \mathbf{y})$. So $T(\mathbf{x}) + T(\mathbf{y})$ is in $T(U)$, and $T(U)$ is closed under vector addition. Let c be any scalar. Then since \mathbf{x} is in U and U is a subspace of V , $c\mathbf{x}$ is in U . Since T

is linear, $T(cx) = cT(x)$ and $cT(x)$ is in $T(U)$. Thus $T(U)$ is closed under scalar multiplication, and $T(U)$ is a subspace of W .

36. Since Z is a subspace of W , $\mathbf{0}_w$ is in Z . Since T is linear, $T(\mathbf{0}_v) = \mathbf{0}_w$. So $\mathbf{0}_v$ is in U . Let x and y be typical elements in U . Then $T(x)$ and $T(y)$ are in Z , and since Z is a subspace of W , $T(x) + T(y)$ is also in Z . Since T is linear, $T(x) + T(y) = T(x + y)$. So $T(x + y)$ is in Z , and $x + y$ is in U . Thus U is closed under vector addition. Let c be any scalar. Then since x is in U , $T(x)$ is in Z . Since Z is a subspace of W , $cT(x)$ is also in Z . Since T is linear, $cT(x) = T(cx)$ and $T(cx)$ is in $T(U)$. Thus cx is in U and U is closed under scalar multiplication. Hence U is a subspace of V .

37. [M] Consider the system with augmented matrix $[A \ w]$. Since

$$[A \ w] = \begin{bmatrix} 1 & 0 & 0 & -1/95 & 1/95 \\ 0 & 1 & 0 & 39/19 & -20/19 \\ 0 & 0 & 1 & 267/95 & -172/95 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

the system is consistent and w is in ColA. Also, since

$$Aw = \begin{bmatrix} 7 & 6 & -4 & 1 \\ -5 & -1 & 0 & -2 \\ 9 & -11 & 7 & -3 \\ 19 & -9 & 7 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -3 \end{bmatrix} = \begin{bmatrix} 14 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

w is not in NulA.

38. [M] Consider the system with augmented matrix $[A \ w]$. Since

$$[A \ w] = \begin{bmatrix} 1 & 0 & -1 & 0 & -2 \\ 0 & 1 & -2 & 0 & -3 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

the system is consistent and w is in ColA. Also, since

$$Aw = \begin{bmatrix} -8 & 5 & -2 & 0 \\ -5 & 2 & 1 & -2 \\ 10 & -8 & 6 & -3 \\ 3 & -2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

w is in NulA.

39. [M]

- a. To show that a_3 and a_5 are in the column space of B , we can row reduce the matrices $[B \ a_3]$ and $[B \ a_5]$:

$$[B \ a_3] = \begin{bmatrix} 1 & 0 & 0 & 1/3 \\ 0 & 1 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$[B \ a_5] = \begin{bmatrix} 1 & 0 & 0 & 10/3 \\ 0 & 1 & 0 & -26/3 \\ 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Since both these systems are consistent, a_3 and a_5 are in the column space of B . Notice that the same conclusions can be drawn by observing the reduced row echelon form for A :

$$A = \begin{bmatrix} 1 & 0 & 1/3 & 0 & 10/3 \\ 0 & 1 & 1/3 & 0 & -26/3 \\ 0 & 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- b. We find the general solution of $Ax = \mathbf{0}$ in terms of the free variables by using the reduced row echelon form of A given above: $x_1 = (-1/3)x_3 - (10/3)x_5$, $x_2 = (-1/3)x_3 + (26/3)x_5$, $x_4 = 4x_5$ with x_3 and x_5 free. So

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_3 \begin{bmatrix} -1/3 \\ -1/3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -10/3 \\ 26/3 \\ 0 \\ 4 \\ 1 \end{bmatrix},$$

and a spanning set for Nul A is

$$\left\{ \begin{bmatrix} -1/3 \\ -1/3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -10/3 \\ 26/3 \\ 0 \\ 4 \\ 1 \end{bmatrix} \right\}.$$

- c. The reduced row echelon form of A shows that the columns of A are linearly dependent and do not span \mathbb{R}^4 . Thus by Theorem 12 in Section 1.9, T is neither one-to-one nor onto.